



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Contents lists available at ScienceDirect

## Diabetes &amp; Metabolic Syndrome: Clinical Research &amp; Reviews

journal homepage: [www.elsevier.com/locate/dsx](http://www.elsevier.com/locate/dsx)

## Original Article

An update on the global vaccine development for coronavirus<sup>☆</sup>Shanay Rab<sup>a</sup>, Afjal<sup>b</sup>, Mohd Javaid<sup>a,\*</sup>, Abid Haleem<sup>a</sup>, Raju Vaishya<sup>c</sup><sup>a</sup> Department of Mechanical Engineering, Jamia Millia Islamia, New Delhi, 110025, India<sup>b</sup> Department of Biotechnology, Amity University, Noida, 201303, India<sup>c</sup> Department of Orthopaedics, Indraprastha Apollo Hospital, Sarita Vihar Mathura Road, New Delhi, 110076, India

## ARTICLE INFO

## Article history:

Received 6 October 2020

Received in revised form

14 October 2020

Accepted 20 October 2020

Dear Editor,

Coronavirus (COVID-19) has created a global crisis and perhaps caused one of the biggest socio-economic tragedies. The world scientific community is struggling hard to develop a solution at the earliest. With the cumulative efforts, the development and implementation of various technological solutions aimed at combating the COVID-19 outbreak are taking shape across the globe. The scientific community is working towards collaborative and fast-paced solutions in terms of therapeutics and vaccines to control the spread of this virus. Globally by September 2020, hundreds of drug companies, biotechnology firms, university research groups, and health organisations were developing more than 500 potential therapies for COVID-19, and they are in various stages of preclinical or clinical research [1–4].

The positive side of early progress towards developing the vaccine for COVID-19 is the evolution of human civilisation with various diseases and previous analysis on SARS-CoV-2, which indicates that SARS-CoV-2 uses the same receptor as SARS-CoV and is approximately 79% genetically similar to SARS-CoV. Various technological platforms for vaccine development are available such as protein subunit vaccines, genetic vaccines, virus vectored vaccines, and monoclonal antibodies for passive immunisation, etc. are under evaluations for COVID-19, with each having their pros and cons [5–7].

For any disease, drug development typically requires several

years to assure the safety & efficacy and approval of regulatory agencies, such as the EMA and the FDA, related to expedite clinically testing procedures. Fig. 1 shows the detailed stage route of any drug development.

As mentioned in Fig. 1, the clinical development stage is crucial for any vaccine development, and it is divided into three different phases. I) Phase 1 trial: The trial vaccine is given to the healthy volunteers, and its safety and dosing are determined. II) Phase 2: The trial establishes an initial reading of efficacy and further explores safety in small numbers of people having the target disease. III) Phase 3: The trial vaccine is given to a large cohort to determine safety and efficacy. One study which has covered clinical research in the 1980–90s, found that only 21.5% of drug candidates that started phase I trials were approved eventually for the manufacturing stage, and during 2006–15, the success rate of obtaining approval from Phase I to successful Phase III trials was under 10% on average, and 16% specifically for vaccines development [8–10]. That is why a successful n-Cov-19 vaccine will require a cautious validation of its efficacy and adverse reactivity to the targeted population.

The World Health Organization (WHO) has implemented an Access to COVID-19 tools accelerator for coordinating global vaccine development program in collaboration with GAVI and the Coalition for Epidemic Preparedness Innovations. As of September 2020, there were 321 companies developing vaccines. However, no company has completed clinical trials to prove its safety and efficacy. In the mid of September 2020, some 42 vaccine candidates were in the clinical development stage in which 33 in Phase III trials and 9 in Phase II-III trials [11].

Russia has become the first country to approve a COVID-19 vaccine. ‘Sputnik V’ – formerly known as the ‘Gam-COVID-Vac’ vaccine, was developed by the Gamaleya Research Institute and Russian Defense Ministry to use the viral vector vaccine method on

<sup>☆</sup> Google scholar link: <https://scholar.google.co.in/citations?user=rfyiwvsAAAAJ&hl=en>.

\* Corresponding author.

E-mail address: [mjavaid@jmi.ac.in](mailto:mjavaid@jmi.ac.in) (M. Javaid).




Fig. 1. The route of vaccine development.

**Table 1**  
Lead developers for COVID-19 vaccine development & trial processes.

Vaccine candidate	Technology used	Lead developers	Clinical trial status	Tentative duration of the trial	References
mRNA-1273	Lipid nanoparticle dispersion containing mRNA	Moderna	Phase III	July 2020–October 2022	[13]
AZD1222	Modified chimp adenovirus vector	The University of Oxford; AstraZeneca; IQVIA; Serum Institute of India	Phase III	May 2020–August 2021	[14,15]
Ad5-nCoV	Recombinant adenovirus type 5 vector	CanSino Biologics; Beijing Institute of Biotechnology of the Academy of Military Medical Sciences	Phase III	March 2020 –December 2021	[16]
CoronaVac	Inactivated vaccine	Sinovac	Phase III	July 2020–October 2021	[17]
JNJ-78436735 (formerly known as Ad26.COV2–S)	Non-replicating viral vector	Johnson & Johnson	Phase III	September 2020 – Early 2021	[18]
BNT162	mRNA	BioNTech, Fosun Pharma, Pfizer	Phase III	April 2020–May 2021	[19]
Yet to decide the name of the vaccine	Inactivated vaccine	Wuhan Institute of Biological Products; China National Pharmaceutical Group (Sinopharm)	Phase III	July 2020–July 2021	[20]

August 11, 2020 [12]. However, the world scientific community has raised doubts about its clinical trials to prove its safety and efficacy. Table 1 shows major COVID-19 vaccine candidates currently in Phase 3 trials. Indeed the list is not comprehensive but a short description of significant work towards vaccine development.

The current global pandemic has crippled the world in terms of development, quality infrastructure, economy, human health, and life. We have provided an up to date overview of the world scientific efforts dedicated to safe and effective vaccine development and possible solution for COVID-19. We believe that the vaccine should be effective and safe for human beings, and hence it should only be permitted for use after duly confirming its efficacy and safety in the clinical trials on a large number of subjects. It should not be rushed into for human use, for political reasons.

**Declaration of competing interest**

There is no conflicts of interest.

**References**

- [1] 4 September Le Tung Thanh, Cramer Jakob P, Chen Robert, Mayhew Stephen. Evolution of the COVID-19 vaccine development landscape. *Nat Rev Drug Discov* 2020. <https://doi.org/10.1038/d41573-020-00151-8>. ISSN 1474-1776. PMID 32887942.
- [2] Draft landscape of COVID 19 candidate vaccines. *World Health Organization*; 17 September 2020. Accessed on . [Accessed 20 September 2020].
- [3] Haleem A, Javaid M, Vaishya R. Effects of COVID 19 pandemic in daily life. *Curr. Med. Res. Pract.* 2020. <https://doi.org/10.1016/j.cmrp.2020.03.011>.
- [4] Rab S, Javaid M, Haleem A, Vaishya R. Face masks are new normal after COVID-19 pandemic. *Diab. Metabol. Syndr.* 2020;14(6):1617–9. <https://doi.org/10.1016/j.dsx.2020.08.021>. Advance online publication.
- [5] Bramanti B, Stenseth NC, Walløe L, Lei X. Plague: a disease which changed the path of human civilization. In: Yang R, Anisimov A, editors. *Yersinia pestis: retrospective and perspective*. Advances in experimental medicine and biology, vol. 918. Dordrecht: Springer; 2016. [https://doi.org/10.1007/978-94-024-0890-4\\_1](https://doi.org/10.1007/978-94-024-0890-4_1).
- [6] Sunita Sajid A, Singh Y, Shukla P. Computational tools for modern vaccine development. *Hum Vaccines Immunother* 2020 3 March;16(3):723–35. <https://doi.org/10.1080/21645515.2019.1670035>. Epub 2019 18 December. PMID: 31545127; PMCID: PMC7227725.
- [7] Graham BS. Advances in antiviral vaccine development. *Immunol Rev* 2013;255(1):230–42. <https://doi.org/10.1111/imr.12098>.

- [8] Cunningham AL, Garçon N, Leo O, Friedland LR, Strugnelli R, Laupèze B, et al. Vaccine development: from concept to early clinical testing. *Vaccine* 2016 Dec 20;34(52):6655–64. <https://doi.org/10.1016/j.vaccine.2016.10.016>. Epub 2016 18 October. PMID: 27769596.
- [9] R&D costs are on the rise. *Med Market Media* 1 June 2003;38(6):14. Archived from the original on 18 October 2016.
- [10] Clinical development success rates: 2006–2015". *BIO Industry Analysis*. June 2016.
- [11] COVID-19 vaccine tracker. *Milken Institute*; 17 September 2020. Retrieved 20 September 2020.
- [12] Safety and immunogenicity of an rAd26 and rAd5 vector-based heterologous prime-boost COVID-19 vaccine in two formulations: two open, non-randomised phase 1/2 studies from Russia, Logunov, Denis Y et al. *Lancet*, Volume 396, Issue 10255, 887 – 897
- [13] Jackson LA, Anderson EJ, Roupael NG, Roberts PC, Makhene M, Coler RN, et al., (mRNA-1273 Study Group). An mRNA vaccine against SARS-CoV-2 – preliminary report. *N Engl J Med* July 2020. <https://doi.org/10.1056/NEJ-Moa2022483>. PMC 7377258.
- [14] A phase III randomized, double-blind, placebo-controlled multicenter study in adults to determine the safety, efficacy, and immunogenicity of AZD1222, a non-replicating ChAdOx1 vector vaccine, for the prevention of COVID-19. *ClinicalTrials.gov (Registry) United States Natl. Libr. Med.* 12 May 2020 NCT04383574. Retrieved, . [Accessed 26 August 2020].
- [15] O'Reilly P. A Phase III study to investigate a vaccine against COVID-19. *ISRCTN* (Registry) 26 May 2020. <https://doi.org/10.1186/ISRCTN89951424>. ISRCTN89951424. Retrieved, . [Accessed 3 August 2020].
- [16] Zhu F, Guan X, Li Y, Huang J, Jiang T, Hou L, et al. Immunogenicity and safety of a recombinant adenovirus type-5-vectored COVID-19 vaccine in healthy adults aged 18 years or older: a randomised, double-blind, placebo-controlled, phase 2 trial". *Lancet* July 2020;396(10249):479–88. [https://doi.org/10.1016/s0140-6736\(20\)31605-6](https://doi.org/10.1016/s0140-6736(20)31605-6). ISSN 0140-6736. PMID 32702299.
- [17] 10 August Zhang Y-J, Zeng G, Pan H-X, Li C-G, others. Immunogenicity and safety of a SARS-CoV-2 inactivated vaccine in healthy adults aged 18–59 years: report of the randomised, double-blind, and placebo-controlled phase 2 clinical trial. *MedRxiv* 2020. <https://doi.org/10.1101/2020.07.31.20161216>. S2CID 221089937.
- [18] Johnson & Johnson initiates pivotal global phase 3 clinical trial of janssen's COVID-19 vaccine candidate. Accessed on, <https://www.jnj.com/johnson-johnson-initiates-pivotal-global-phase-3-clinical-trial-of-janssens-covid-19-vaccine-candidate>. [Accessed 26 September 2020].
- [19] Sahin U, Muik A, Derhovanessian E, Vogler I, Kranz LM, Vormehr M, et al. Concurrent human antibody and TH1 type T-cell responses elicited by a COVID-19 RNA vaccine. *MedRxiv* (Preprint) 2020. <https://doi.org/10.1101/2020.07.17.20140533>.
- [20] Xia S, Duan K, Zhang Y, Zhao D, et al. Effect of an inactivated vaccine against SARS-CoV-2 on safety and immunogenicity outcomes: interim analysis of 2 randomised clinical trials. *J Am Med Assoc* 13 August 2020. <https://doi.org/10.1001/jama.2020.15543>. PMC 7426884. PMID 32789505.